

# Proxalutamide for COVID-19: real-time meta analysis of 4 studies

@CovidAnalysis, March 2024, Version 11  
<https://c19early.org/xmeta.html>

## Abstract

Statistically significant lower risk is seen for mortality, ventilation, hospitalization, recovery, and viral clearance. 4 studies from 2 independent teams in 2 countries show statistically significant improvements.



Meta analysis using the most serious outcome reported shows 78% [70-83%] lower risk. Results are similar for peer-reviewed studies. Currently all studies are RCTs.

Studies to date are from only 2 different groups.

No treatment or intervention is 100% effective. All practical, effective, and safe means should be used based on risk/benefit analysis. Multiple treatments are typically used in combination, and other treatments may be more effective.

All data to reproduce this paper and sources are in the appendix.

Zheng present another meta analysis for proxalutamide, showing significant improvements for mortality, mechanical ventilation, hospitalization, and clinical improvement.



## HIGHLIGHTS


Proxalutamide reduces risk for COVID-19 with very high confidence for mortality and in pooled analysis, high confidence for hospitalization, and low confidence for ventilation, recovery, and viral clearance.

Proxalutamide was the 30th treatment shown effective with  $\geq 3$  clinical studies in December 2021, now known with  $p < 0.0000000001$  from 4 studies, and recognized in 3 countries.

We show traditional outcome specific analyses and combined evidence from all studies, incorporating treatment delay, a primary confounding factor in COVID-19 studies.

Real-time updates and corrections, transparent analysis with all results in the same format, consistent protocol for 66 treatments.

4 proxalutamide COVID-19 studies




A

Efficacy in COVID-19 proxalutamide studies (pooled effects) c19early.org  
March 2024




B

Efficacy in COVID-19 studies (pooled effects) c19early.org  
March 2024



C

## Timeline of COVID-19 proxalutamide studies (pooled effects)



**Figure 1. A. Random effects meta-analysis.** This plot shows pooled effects, see the specific outcome analyses for individual outcomes, and the heterogeneity section for discussion. Effect extraction is pre-specified, using the most serious outcome reported. For details of effect extraction see the [appendix](#). **B. Scatter plot showing the most serious outcome in all studies, and for studies within each stage.** Diamonds shows the results of random effects meta-analysis. **C. Results within the context of multiple COVID-19 treatments.** 0.6% of 6,686 proposed treatments show efficacy [c19early.org](#). **D. Timeline of results in proxalutamide studies.** The marked dates indicate the time when efficacy was known with a statistically significant improvement of  $\geq 10\%$  from  $\geq 3$  studies for pooled outcomes, one or more specific outcome, pooled outcomes in RCTs, and one or more specific outcome in RCTs.


## Introduction

**Immediate treatment recommended.** SARS-CoV-2 infection primarily begins in the upper respiratory tract and may progress to the lower respiratory tract, other tissues, and the nervous and cardiovascular systems, which may lead to cytokine storm, pneumonia, ARDS, neurological issues [Scardua-Silva, Yang](#), cardiovascular complications [Eberhardt](#), organ failure, and death. Minimizing replication as early as possible is recommended.

**Many treatments are expected to modulate infection.** SARS-CoV-2 infection and replication involves the complex interplay of 50+ host and viral proteins and other factors [Note A, Malone, Murigneux, Lv, Lui](#), providing many therapeutic targets for which many existing compounds have known activity. Scientists have predicted that over 6,000 compounds may reduce COVID-19 risk [c19early.org](#) (B), either by directly minimizing infection or replication, by supporting immune system function, or by minimizing secondary complications.

**Analysis.** We analyze all significant controlled studies of proxalutamide for COVID-19. Search methods, inclusion criteria, effect extraction criteria (more serious outcomes have priority), all individual study data, PRISMA answers, and statistical methods are detailed in Appendix 1. We present random effects meta-analysis results for all studies, studies within each treatment stage, individual outcomes, peer-reviewed studies, and Randomized Controlled Trials (RCTs).

**Treatment timing.** Figure 2 shows stages of possible treatment for COVID-19. Prophylaxis refers to regularly taking medication before becoming sick, in order to prevent or minimize infection. Early Treatment refers to treatment immediately or soon after symptoms appear, while Late Treatment refers to more delayed treatment.



**Figure 2.** Treatment stages.

## Preclinical Research

2 *In Vitro* studies support the efficacy of proxalutamide *Ma, Qiao*.

Preclinical research is an important part of the development of treatments, however results may be very different in clinical trials. Preclinical results are not used in this paper.

## Results

Table 1 summarizes the results for all stages combined, for Randomized Controlled Trials, for peer-reviewed studies, and for specific outcomes. Table 2 shows results by treatment stage. Figure 3, 4, 5, 6, 7, 8, and 9 show forest plots for random effects meta-analysis of all studies with pooled effects, mortality results, ventilation, hospitalization, recovery, viral clearance, and peer reviewed studies.

	<i>Improvement</i>	<i>Studies</i>	<i>Patients</i>	<i>Authors</i>
All studies	78% [70-83%] ****	4	1,953	38
Peer-reviewed studies	78% [70-84%] ****	2	1,046	30
Randomized Controlled Trials	78% [70-83%] ****	4	1,953	38
Mortality	78% [70-83%] ****	4	1,953	38
Ventilation	95% [60-99%] **	2	445	22
Hospitalization	72% [19-90%] *	4	1,953	38
RCT mortality	78% [70-83%] ****	4	1,953	38
RCT hospitalization	72% [19-90%] *	4	1,953	38

**Table 1.** Random effects meta-analysis for all stages combined, for Randomized Controlled Trials, for peer-reviewed studies, and for specific outcomes. Results show the percentage improvement with treatment and the 95% confidence interval.

\*\*  $p < 0.01$  \*\*\*  $p < 0.001$  \*\*\*\*  $p < 0.0001$ .



## 4 proxalutamide COVID-19 mortality results

c19early.org

March 2024




Figure 4. Random effects meta-analysis for mortality results.

## 2 proxalutamide COVID-19 mechanical ventilation results

c19early.org

March 2024




Figure 5. Random effects meta-analysis for ventilation.

## 4 proxalutamide COVID-19 hospitalization results

c19early.org

March 2024





Figure 6. Random effects meta-analysis for hospitalization.

## 1 proxalutamide COVID-19 recovery result


c19early.org  
March 2024



**Figure 7.** Random effects meta-analysis for recovery.

## 1 proxalutamide COVID-19 viral clearance result


c19early.org  
March 2024



**Figure 8.** Random effects meta-analysis for viral clearance.

## 2 proxalutamide COVID-19 peer reviewed studies

c19early.org  
March 2024



**Figure 9.** Random effects meta-analysis for peer reviewed studies. Effect extraction is pre-specified, using the most serious outcome reported, see the appendix for details. Zeraatkar *et al.* analyze 356 COVID-19 trials, finding no significant evidence that preprint results are inconsistent with peer-reviewed studies. They also show extremely long peer-review delays, with a median of 6 months to journal publication. A six month delay was equivalent to around 1.5 million deaths during the first two years of the pandemic. Authors recommend using preprint evidence, with appropriate checks for potential falsified data, which provides higher certainty much earlier. Davidson *et al.* also showed no important difference between meta analysis results of preprints and peer-reviewed publications for COVID-19, based on 37 meta analyses including 114 trials.

# Randomized Controlled Trials (RCTs)

Currently all studies are RCTs.

## Heterogeneity

Heterogeneity in COVID-19 studies arises from many factors including:


**Treatment delay.** The time between infection or the onset of symptoms and treatment may critically affect how well a treatment works. For example an antiviral may be very effective when used early but may not be effective in late stage disease, and may even be harmful. Oseltamivir, for example, is generally only considered effective for influenza when used within 0-36 or 0-48 hours *McLean, Treanor*. Baloxavir studies for influenza also show that treatment delay is critical — *Ikematsu* report an 86% reduction in cases for post-exposure prophylaxis, *Hayden* show a 33 hour reduction in the time to alleviation of symptoms for treatment within 24 hours and a reduction of 13 hours for treatment within 24-48 hours, and *Kumar* report only 2.5 hours improvement for inpatient treatment.

<i>Treatment delay</i>	<i>Result</i>
Post exposure prophylaxis	<b>86% fewer cases</b> <i>Ikematsu</i>
<24 hours	<b>-33 hours symptoms</b> <i>Hayden</i>
24-48 hours	<b>-13 hours symptoms</b> <i>Hayden</i>
Inpatients	<b>-2.5 hours to improvement</b> <i>Kumar</i>

**Table 3.** Studies of baloxavir for influenza show that early treatment is more effective.

Figure 10 shows a mixed-effects meta-regression for efficacy as a function of treatment delay in COVID-19 studies from 66 treatments, showing that efficacy declines rapidly with treatment delay. Early treatment is critical for COVID-19.





**Figure 10. Early treatment is more effective.** Meta-regression showing efficacy as a function of treatment delay in COVID-19 studies from 66 treatments.

**Patient demographics.** Details of the patient population including age and comorbidities may critically affect how well a treatment works. For example, many COVID-19 studies with relatively young low-comorbidity patients show all patients recovering quickly with or without treatment. In such cases, there is little room for an effective treatment to improve results (as in *López-Medina*).

**Effect measured.** Efficacy may differ significantly depending on the effect measured, for example a treatment may be very effective at reducing mortality, but less effective at minimizing cases or hospitalization. Or a treatment may have no effect on viral clearance while still being effective at reducing mortality.

**Variants.** There are many different variants of SARS-CoV-2 and efficacy may depend critically on the distribution of variants encountered by the patients in a study. For example, the Gamma variant shows significantly different characteristics *Faria, Karita, Nonaka, Zavascki*. Different mechanisms of action may be more or less effective depending on variants, for example the viral entry process for the omicron variant has moved towards TMPRSS2-independent fusion, suggesting that TMPRSS2 inhibitors may be less effective *Peacock, Willett*.

**Regimen.** Effectiveness may depend strongly on the dosage and treatment regimen.

**Other treatments.** The use of other treatments may significantly affect outcomes, including anything from supplements, other medications, or other kinds of treatment such as prone positioning.


**Medication quality.** The quality of medications may vary significantly between manufacturers and production batches, which may significantly affect efficacy and safety. *Williams* analyze ivermectin from 11 different sources, showing highly variable antiparasitic efficacy across different manufacturers. *Xu* analyze a treatment from two different manufacturers, showing 9 different impurities, with significantly different concentrations for each manufacturer.

**Pooled outcome analysis.** We present both pooled analyses and specific outcome analyses. Notably, pooled analysis often results in earlier detection of efficacy as shown in Figure 11. For many COVID-19 treatments, a reduction in mortality logically follows from a reduction in hospitalization, which follows from a reduction in symptomatic cases, etc. An antiviral tested with a low-risk population may report zero mortality in both arms, however a reduction in severity and improved viral clearance may translate into lower mortality among a high-risk population, and including these results in pooled analysis allows faster detection of efficacy. Trials with high-risk patients may also be restricted due to ethical concerns for treatments that are known or expected to be effective.

Pooled analysis enables using more of the available information. While there is much more information available, for example dose-response relationships, the advantage of the method used here is simplicity and transparency. Note that pooled analysis could hide efficacy, for example a treatment that is beneficial for late stage patients but has no effect on viral replication or early stage disease could show no efficacy in pooled analysis if most studies only examine viral clearance. While we present pooled results, we also present individual outcome analyses, which may be more informative for specific use cases.

**Pooled outcomes identify efficacy faster.** Currently, 44 of the treatments we analyze show statistically significant efficacy or harm, defined as  $\geq 10\%$  decreased risk or  $>0\%$  increased risk from  $\geq 3$  studies. 88% of treatments showing statistically significant efficacy/harm with pooled effects have been confirmed with one or more specific outcomes, with a mean delay of 3.6 months. When restricting to RCTs only, 50% of treatments showing statistically significant efficacy/harm with pooled effects have been confirmed with one or more specific outcomes, with a mean delay of 6.1 months.

## Time when COVID-19 studies showed efficacy



**Figure 11.** The time when studies showed that treatments were effective, defined as statistically significant improvement of  $\geq 10\%$  from  $\geq 3$  studies. Pooled results typically show efficacy earlier than specific outcome results. Results from all studies often shows efficacy much earlier than when restricting to RCTs. Results reflect conditions as used in trials to date, these depend on the population treated, treatment delay, and treatment regimen.


**Meta analysis.** The distribution of studies will alter the outcome of a meta analysis. Consider a simplified example where everything is equal except for the treatment delay, and effectiveness decreases to zero or below with increasing delay. If there are many studies using very late treatment, the outcome may be negative, even though early treatment is very effective. This may have a greater effect than pooling different outcomes such as mortality and hospitalization. For example a treatment may have 50% efficacy for mortality but only 40% for hospitalization when used within 48 hours. However efficacy could be 0% when used late.

All meta analyses combine heterogeneous studies, varying in population, variants, and potentially all factors above, and therefore may obscure efficacy by including studies where treatment is less effective. Generally, we expect the estimated effect size from meta analysis to be less than that for the optimal case. Looking at all studies is valuable for providing an overview of all research, important to avoid cherry-picking, and informative when a positive result is found despite combining less-optimal situations. However, the resulting estimate does not apply to specific cases such as early treatment in high-risk populations. While we present results for all studies, we also present treatment time and individual outcome analyses, which may be more informative for specific use cases.

## Discussion

**Publication bias.** Publishing is often biased towards positive results. Trials with patented drugs may have a financial conflict of interest that results in positive studies being more likely to be published, or bias towards more positive results. For example with molnupiravir, trials with negative results remain unpublished to date (CTRI/2021/05/033864 and CTRI/2021/08/0354242). For proxalutamide, there is currently not enough data to evaluate publication bias with high confidence.

**Funnel plot analysis.** Funnel plots have traditionally been used for analyzing publication bias. This is invalid for COVID-19 acute treatment trials — the underlying assumptions are invalid, which we can demonstrate with a simple example. Consider a set of hypothetical perfect trials with no bias. Figure 12 plot A shows a funnel plot for a simulation of 80 perfect trials, with random group sizes, and each patient's outcome randomly sampled (10% control event probability, and a 30% effect size for treatment). Analysis shows no asymmetry ( $p > 0.05$ ). In plot B, we add a single typical variation in COVID-19 treatment trials — treatment delay. Consider that efficacy varies from 90% for treatment within 24 hours, reducing to 10% when treatment is delayed 3 days. In plot B, each trial's treatment delay is randomly selected. Analysis now shows highly significant asymmetry,  $p < 0.0001$ , with six variants of Egger's test all showing  $p < 0.05$  *Egger, Harbord, Macaskill, Moreno, Peters, Rothstein, Rücker, Stanley*. Note that these tests fail even though treatment delay is uniformly distributed. In reality treatment delay is more complex — each trial has a different distribution of delays across patients, and the distribution across trials may be biased (e.g., late treatment trials may be more common). Similarly, many other variations in trials may produce asymmetry, including dose, administration, duration of treatment, differences in SOC, comorbidities, age, variants, and bias in design, implementation, analysis, and reporting.



**Figure 12.** Example funnel plot analysis for simulated perfect trials.

**Limitations.** Summary statistics from meta analysis necessarily lose information. As with all meta analyses, studies are heterogeneous, with differences in treatment delay, treatment regimen, patient demographics, variants, conflicts of interest, standard of care, and other factors. We provide analyses by specific outcomes and by treatment delay, and we aim to identify key characteristics in the forest plots and summaries. Results should be viewed in the context of study characteristics.

Some analyses classify treatment based on early or late administration, as done here, while others distinguish between mild, moderate, and severe cases. Viral load does not indicate degree of symptoms — for example patients may have a high viral load while being asymptomatic. With regard to treatments that have antiviral properties, timing of treatment is critical — late administration may be less helpful regardless of severity.

Details of treatment delay per patient is often not available. For example, a study may treat 90% of patients relatively early, but the events driving the outcome may come from 10% of patients treated very late. Our 5 day cutoff for early treatment may be too conservative, 5 days may be too late in many cases.

Comparison across treatments is confounded by differences in the studies performed, for example dose, variants, and conflicts of interest. Trials affiliated with special interests may use designs better suited to the preferred outcome.

In some cases, the most serious outcome has very few events, resulting in lower confidence results being used in pooled analysis, however the method is simpler and more transparent. This is less critical as the number of studies increases. Restriction to outcomes with sufficient power may be beneficial in pooled analysis and improve accuracy when there are few studies, however we maintain our pre-specified method to avoid any retrospective changes.

Studies show that combinations of treatments can be highly synergistic and may result in many times greater efficacy than individual treatments alone *Alsaiddi, Andreani, De Forni, Fiaschi, Jeffreys, Jitobaom, Jitobaom (B), Ostrov, Said, Thairu, Wan*. Therefore standard of care may be critical and benefits may diminish or disappear if standard of care does not include certain treatments.

This real-time analysis is constantly updated based on submissions. Accuracy benefits from widespread review and submission of updates and corrections from reviewers. Less popular treatments may receive fewer reviews.

No treatment, vaccine, or intervention is 100% available and effective for all current and future variants. Efficacy may vary significantly with different variants and within different populations. All treatments have potential side effects. Propensity to experience side effects may be predicted in advance by qualified physicians. We do not provide medical advice. Before taking any medication, consult a qualified physician who can compare all options, provide personalized advice, and provide details of risks and benefits based on individual medical history and situations.

**Notes.** *Zheng* present another meta analysis for proxalutamide, showing significant improvements for mortality, mechanical ventilation, hospitalization, and clinical improvement.

## Conclusion


Studies to date show that proxalutamide is an effective treatment for COVID-19. Statistically significant lower risk is seen for mortality, ventilation, hospitalization, recovery, and viral clearance. 4 studies from 2 independent teams in 2 countries show statistically significant improvements. Meta analysis using the most serious outcome reported shows 78% [70-83%] lower risk. Results are similar for peer-reviewed studies. Currently all studies are RCTs.

Studies to date are from only 2 different groups.

*Zheng* present another meta analysis for proxalutamide, showing significant improvements for mortality, mechanical ventilation, hospitalization, and clinical improvement.

## Study Notes

### Cadegiani




#### SEE ALSO

The High-Impact Medical Journal Editors Harassment Of The World's Leading Clinical Researcher of Repurposed Dr...

*Cadegiani*: RCT 177 women in Brazil, 75 treated with proxalutamide, showing significantly lower hospitalization with treatment.

### Cadegiani




#### SEE ALSO

The High-Impact Medical Journal Editors Harassment Of The World's Leading Clinical Researcher of Repurposed Dr...


*Cadegiani (B)*: RCT 778 hospitalized patients in Brazil, 423 treated with proxalutamide, showing significantly lower mortality and improved recovery with treatment. NCT04728802 and NCT05126628. Authors note that cases in this trial were predominantly the P.1 Gamma variant, for which proxalutamide may be more effective compared to other variants.

## Kintor



*Kintor*: RCT 733 outpatients, 99% in the USA, showing lower hospitalization/death, and significantly reduced viral load with proxalutamide treatment. The viral clearance result is from *Ma et al.*.

## McCoy



### SEE ALSO

The High-Impact Medical Journal Editors Harassment Of The World's Leading Clinical Researcher of Repurposed Dr...

*McCoy*: RCT 268 male patients in Brazil, 134 treated with proxalutamide, showing significantly lower hospitalization and mechanical ventilation. NCT04446429.

This paper was censored without details or author response, and the editors have ignored the authors, see [twitter.com](https://twitter.com).

## Appendix 1. Methods and Data

We perform ongoing searches of PubMed, medRxiv, Europe PMC, ClinicalTrials.gov, The Cochrane Library, Google Scholar, Research Square, ScienceDirect, Oxford University Press, the reference lists of other studies and meta-analyses, and submissions to the site c19early.org. Search terms are proxalutamide and COVID-19 or SARS-CoV-2.

Automated searches are performed twice daily, with all matches reviewed for inclusion. All studies regarding the use of proxalutamide for COVID-19 that report a comparison with a control group are included in the main analysis. This is a living analysis and is updated regularly.

We extracted effect sizes and associated data from all studies. If studies report multiple kinds of effects then the most serious outcome is used in pooled analysis, while other outcomes are included in the outcome specific analyses. For example, if effects for mortality and cases are both reported, the effect for mortality is used, this may be different to the effect that a study focused on. If symptomatic results are reported at multiple times, we used the latest time, for example if mortality results are provided at 14 days and 28 days, the results at 28 days have preference. Mortality alone is preferred over combined outcomes. Outcomes with zero events in both arms are not used, the next most serious outcome with one or more events is used. For example, in low-risk populations with no mortality, a reduction in mortality with treatment is not possible, however a reduction in hospitalization, for example, is still valuable. Clinical outcomes are considered more important than viral test status. When basically all patients recover in both treatment and control groups, preference for viral clearance and recovery is given to results mid-recovery where available. After most or all patients have recovered there is little or no room for an effective treatment to do better, however faster recovery is valuable. If only individual symptom data is available, the most serious symptom has priority, for example difficulty breathing or low SpO<sub>2</sub> is more important than cough. When results provide an odds ratio, we compute the relative risk when possible, or convert to a relative risk according to *Zhang*. Reported confidence intervals and *p*-values were used when available, using adjusted values when provided. If multiple types of adjustments are reported propensity score matching and multivariable regression has preference over propensity score matching or weighting, which has preference over multivariable regression. Adjusted results have preference over unadjusted results for a more serious outcome when the adjustments significantly alter results. When needed, conversion between reported *p*-values and confidence intervals followed *Altman, Altman (B)*, and Fisher's exact test was used to calculate *p*-values for event data. If continuity correction for zero values is required, we use the reciprocal of the opposite arm with the sum of the correction factors equal to 1 *Sweeting*. Results are expressed with RR < 1.0 favoring treatment, and using the risk of a negative outcome when applicable (for example, the risk of death rather than the risk of survival). If studies only report relative continuous values such as relative times, the ratio of the time for the treatment group versus the time for the control group is used. Calculations are done in Python (3.12.2) with scipy (1.12.0), pythonmeta (1.26), numpy (1.26.4), statsmodels (0.14.1), and plotly (5.19.0).

Forest plots are computed using PythonMeta *Deng* with the DerSimonian and Laird random effects model (the fixed effect assumption is not plausible in this case) and inverse variance weighting. Results are presented with 95% confidence intervals. Heterogeneity among studies was assessed using the I<sup>2</sup> statistic. Mixed-effects meta-regression results are computed with R (4.1.2) using the metafor (3.0-2) and rms (6.2-0) packages, and using the most serious sufficiently powered outcome. For all statistical tests, a *p*-value less than 0.05 was considered statistically significant. Grobid 0.8.0 is used to parse PDF documents.

We have classified studies as early treatment if most patients are not already at a severe stage at the time of treatment (for example based on oxygen status or lung involvement), and treatment started within 5 days of the onset of symptoms. If studies contain a mix of early treatment and late treatment patients, we consider the treatment time of patients contributing most to the events (for example, consider a study where most patients are treated early but late treatment patients are included, and all mortality events were observed with late treatment patients). We note that a shorter time may be preferable. Antivirals are typically only considered effective when used within a shorter timeframe, for example 0-36 or 0-48 hours for oseltamivir, with longer delays not being effective *McLean, Treanor*.

We received no funding, this research is done in our spare time. We have no affiliations with any pharmaceutical companies or political parties.

A summary of study results is below. Please submit updates and corrections at <https://c19early.org/xmeta.html>.

## Early treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. For pooled analyses, the first (most serious) outcome is used, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.



<p><i>Cadegiani</i>, 7/10/2021, Double Blind Randomized Controlled Trial, Brazil, preprint, 7 authors, study period 4 January, 2021 - 28 February, 2021.</p>	<p>risk of death, 63.4% lower, RR 0.37, <math>p = 1.00</math>, treatment 0 of 75 (0.0%), control 1 of 102 (1.0%), NNT 102, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).</p>
	<p>risk of mechanical ventilation, 89.7% lower, RR 0.10, <math>p = 0.07</math>, treatment 0 of 75 (0.0%), control 5 of 102 (4.9%), NNT 20, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).</p>
	<p>risk of hospitalization, 85.7% lower, RR 0.14, <math>p &lt; 0.001</math>, treatment 2 of 75 (2.7%), control 19 of 102 (18.6%), NNT 6.3.</p>
<p><i>Kintor</i>, 4/5/2022, Double Blind Randomized Controlled Trial, placebo-controlled, USA, preprint, 1 author, study period 5 March, 2021 - 1 April, 2022, trial NCT04870606 (history).</p>	<p>risk of death, 66.7% lower, RR 0.33, <math>p = 1.00</math>, treatment 0 of 365 (0.0%), control 1 of 365 (0.3%), NNT 365, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), 1+ days of treatment, group size approximated.</p>
	<p>risk of hospitalization, 50.0% lower, RR 0.50, <math>p = 0.38</math>, treatment 4 of 365 (1.1%), control 8 of 365 (2.2%), NNT 91, 1+ days of treatment, group size approximated.</p>
	<p>risk of death, 66.6% lower, RR 0.33, <math>p = 1.00</math>, treatment 0 of 360 (0.0%), control 1 of 361 (0.3%), NNT 361, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), &gt;1 day of treatment, group size approximated.</p>
	<p>risk of hospitalization, 71.3% lower, RR 0.29, <math>p = 0.18</math>, treatment 2 of 360 (0.6%), control 7 of 361 (1.9%), NNT 72, &gt;1 day of treatment, group size approximated.</p>
	<p>risk of death, 66.6% lower, RR 0.33, <math>p = 1.00</math>, treatment 0 of 346 (0.0%), control 1 of 347 (0.3%), NNT 347, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), &gt;7 days of treatment, group size approximated.</p>
	<p>risk of hospitalization, 92.3% lower, RR 0.08, <math>p = 0.03</math>, treatment 0 of 346 (0.0%), control 6 of 347 (1.7%), NNT 58, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm), &gt;7 days of treatment, group size approximated.</p>
	<p>risk of no viral clearance, 73.9% lower, RR 0.26, <math>p &lt; 0.001</math>, day 7.</p>
<p><i>McCoy</i>, 12/30/2020, Double Blind Randomized Controlled Trial, Brazil, peer-reviewed, 15 authors, study period 15 June, 2020 - 28 July, 2020, trial NCT04446429 (history).</p>	<p>risk of death, 80.0% lower, RR 0.20, <math>p = 0.50</math>, treatment 0 of 134 (0.0%), control 2 of 134 (1.5%), NNT 67, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).</p>
	<p>risk of mechanical ventilation, 97.1% lower, RR 0.03, <math>p &lt; 0.001</math>, treatment 0 of 134 (0.0%), control 17 of 134 (12.7%), NNT 7.9, relative risk is not 0 because of continuity correction due to zero events (with reciprocal of the contrasting arm).</p>



<p>risk of hospitalization, 91.0% lower, RR 0.09, <math>p &lt; 0.001</math>, treatment 3 of 134 (2.2%), control 35 of 134 (26.1%), NNT 4.2.</p>
---

## Late treatment

Effect extraction follows pre-specified rules as detailed above and gives priority to more serious outcomes. For pooled analyses, the first (most serious) outcome is used, which may differ from the effect a paper focuses on. Other outcomes are used in outcome specific analyses.

<p><i>Cadegiani (B)</i>, 12/25/2021, Double Blind Randomized Controlled Trial, Brazil, peer-reviewed, 15 authors, study period 1 February, 2021 - 15 April, 2021, trial NCT04728802 (history).</p>	<p>risk of death, 78.0% lower, RR 0.22, <math>p &lt; 0.001</math>, treatment 45 of 423 (10.6%), control 171 of 355 (48.2%), NNT 2.7, adjusted per study, 28 days, Cox proportional hazards.</p>
	<p>risk of death, 79.0% lower, RR 0.21, <math>p &lt; 0.001</math>, treatment 34 of 423 (8.0%), control 138 of 355 (38.9%), NNT 3.2, adjusted per study, 14 days, Cox proportional hazards.</p>
	<p>recovery rate, RR 0.55, <math>p &lt; 0.001</math>, treatment 423, control 355, adjusted per study, inverted to make <math>RR &lt; 1</math> favor treatment, 28 days, Cox proportional hazards.</p>
	<p>recovery rate, RR 0.45, <math>p &lt; 0.001</math>, treatment 423, control 355, adjusted per study, inverted to make <math>RR &lt; 1</math> favor treatment, 14 days, Cox proportional hazards, primary outcome.</p>
	<p>hospitalization time, 33.3% lower, relative time 0.67, <math>p &lt; 0.001</math>, treatment 423, control 355.</p>

## Supplementary Data

Supplementary Data

## Footnotes

- a. Viral infection and replication involves attachment, entry, uncoating and release, genome replication and transcription, translation and protein processing, assembly and budding, and release. Each step can be disrupted by therapeutics.

## References

1. **Alsaidi** et al., *Griffithsin and Carrageenan Combination Results in Antiviral Synergy against SARS-CoV-1 and 2 in a Pseudoviral Model*, Marine Drugs, doi:10.3390/md19080418.
2. **Altman**, D., *How to obtain the P value from a confidence interval*, BMJ, doi:10.1136/bmj.d2304.
3. **Altman (B)** et al., *How to obtain the confidence interval from a P value*, BMJ, doi:10.1136/bmj.d2090.
4. **Andreani** et al., *In vitro testing of combined hydroxychloroquine and azithromycin on SARS-CoV-2 shows synergistic effect*, Microbial Pathogenesis, doi:10.1016/j.micpath.2020.104228.
5. **c19early.org**, c19early.org/timeline.html.

6. **c19early.org (B)**, [c19early.org/treatments.html](https://c19early.org/treatments.html).
7. **Cadegiani** et al., *Proxalutamide (GT0918) Reduces the Rate of Hospitalization in mild-to-moderate COVID-19 Female Patients: A Randomized Double-Blinded Placebo-Controlled Two-Arm Parallel Trial*, medRxiv, doi:10.1101/2021.07.06.21260086.
8. **Cadegiani (B)** et al., *Final Results of a Randomized, Placebo-Controlled, Two-Arm, Parallel Clinical Trial of Proxalutamide for Hospitalized COVID-19 Patients: A Multiregional, Joint Analysis of the Proxa-Rescue AndroCoV Trial*, Cureus, doi:10.7759/cureus.20691.
9. **Davidson** et al., *No evidence of important difference in summary treatment effects between COVID-19 preprints and peer-reviewed publications: a meta-epidemiological study*, Journal of Clinical Epidemiology, doi:10.1016/j.jclinepi.2023.08.011.
10. **De Forni** et al., *Synergistic drug combinations designed to fully suppress SARS-CoV-2 in the lung of COVID-19 patients*, PLoS ONE, doi:10.1371/journal.pone.0276751.
11. **Deng**, H., *PyMeta, Python module for meta-analysis*, [www.pymeta.com/](http://www.pymeta.com/).
12. **Eberhardt** et al., *SARS-CoV-2 infection triggers pro-atherogenic inflammatory responses in human coronary vessels*, Nature Cardiovascular Research, doi:10.1038/s44161-023-00336-5.
13. **Egger** et al., *Bias in meta-analysis detected by a simple, graphical test*, BMJ, doi:10.1136/bmj.315.7109.629.
14. **Faria** et al., *Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus, Brazil*, Science, doi:10.1126/science.abh2644.
15. **Fiaschi** et al., *In Vitro Combinatorial Activity of Direct Acting Antivirals and Monoclonal Antibodies against the Ancestral B.1 and BQ.1.1 SARS-CoV-2 Viral Variants*, Viruses, doi:10.3390/v16020168.
16. **Harbord** et al., *A modified test for small-study effects in meta-analyses of controlled trials with binary endpoints*, Statistics in Medicine, doi:10.1002/sim.2380.
17. **Hayden** et al., *Baloxavir Marboxil for Uncomplicated Influenza in Adults and Adolescents*, New England Journal of Medicine, doi:10.1056/NEJMoa1716197.
18. **Ikematsu** et al., *Baloxavir Marboxil for Prophylaxis against Influenza in Household Contacts*, New England Journal of Medicine, doi:10.1056/NEJMoa1915341.
19. **Jeffreys** et al., *Remdesivir-ivermectin combination displays synergistic interaction with improved in vitro activity against SARS-CoV-2*, International Journal of Antimicrobial Agents, doi:10.1016/j.ijantimicag.2022.106542.
20. **Jitobaom** et al., *Favipiravir and Ivermectin Showed in Vitro Synergistic Antiviral Activity against SARS-CoV-2*, Research Square, doi:10.21203/rs.3.rs-941811/v1.
21. **Jitobaom (B)** et al., *Synergistic anti-SARS-CoV-2 activity of repurposed anti-parasitic drug combinations*, BMC Pharmacology and Toxicology, doi:10.1186/s40360-022-00580-8.
22. **Karita** et al., *Trajectory of viral load in a prospective population-based cohort with incident SARS-CoV-2 G614 infection*, medRxiv, doi:10.1101/2021.08.27.21262754.
23. **Kintor**, *Kintor Pharma's Proxalutamide Demonstrated Reduction in Hospitalization/Mortality for Patients with Mild to Moderate COVID-19 in Phase III MRCT Study*, Press Release, [www.prnewswire.com/news-releases/kintor-pharmas-proxalutamide-demonstrated-reduction-in-hospitalizationmortality-for-patients-with-mild-to-moderate-covid-19-in-phase-iii-mrct-study-301518525.html](https://www.prnewswire.com/news-releases/kintor-pharmas-proxalutamide-demonstrated-reduction-in-hospitalizationmortality-for-patients-with-mild-to-moderate-covid-19-in-phase-iii-mrct-study-301518525.html).
24. **Kumar** et al., *Combining baloxavir marboxil with standard-of-care neuraminidase inhibitor in patients hospitalised with severe influenza (FLAGSTONE): a randomised, parallel-group, double-blind, placebo-controlled, superiority trial*, The Lancet Infectious Diseases, doi:10.1016/S1473-3099(21)00469-2.
25. **López-Medina** et al., *Effect of Ivermectin on Time to Resolution of Symptoms Among Adults With Mild COVID-19: A Randomized Clinical Trial*, JAMA, doi:10.1001/jama.2021.3071.
26. **Lui** et al., *Nsp1 facilitates SARS-CoV-2 replication through calcineurin-NFAT signaling*, Virology, doi:10.1128/mbio.00392-24.
27. **Lv** et al., *Host proviral and antiviral factors for SARS-CoV-2*, Virus Genes, doi:10.1007/s11262-021-01869-2.

28. **Ma** et al., *Mechanisms of action (MOA) for proxalutamide, an androgen receptor (AR) antagonist, for the treatment of mild, moderate and severe COVID-19 patients*, AACR Annual Meeting 2022, [www.abstractsonline.com/pp8/#!/10517/presentation/18533](http://www.abstractsonline.com/pp8/#!/10517/presentation/18533).
29. **Macaskill** et al., *A comparison of methods to detect publication bias in meta-analysis*, *Statistics in Medicine*, doi:10.1002/sim.698.
30. **Malone** et al., *Structures and functions of coronavirus replication–transcription complexes and their relevance for SARS-CoV-2 drug design*, *Nature Reviews Molecular Cell Biology*, doi:10.1038/s41580-021-00432-z.
31. **McCoy** et al., *Proxalutamide (GT0918) Reduces the Rate of Hospitalization for COVID-19 Male Outpatients: A Randomized Double-Blinded Placebo-Controlled Trial*, *Frontiers in Medicine*, doi:10.3389/fmed.2021.668698.
32. **McLean** et al., *Impact of Late Oseltamivir Treatment on Influenza Symptoms in the Outpatient Setting: Results of a Randomized Trial*, *Open Forum Infect. Dis.* September 2015, 2:3, doi:10.1093/ofid/ofv100.
33. **Moreno** et al., *Assessment of regression-based methods to adjust for publication bias through a comprehensive simulation study*, *BMC Medical Research Methodology*, doi:10.1186/1471-2288-9-2.
34. **Murigneux** et al., *Proteomic analysis of SARS-CoV-2 particles unveils a key role of G3BP proteins in viral assembly*, *Nature Communications*, doi:10.1038/s41467-024-44958-0.
35. **Nonaka** et al., *SARS-CoV-2 variant of concern P.1 (Gamma) infection in young and middle-aged patients admitted to the intensive care units of a single hospital in Salvador, Northeast Brazil, February 2021*, *International Journal of Infectious Diseases*, doi:10.1016/j.ijid.2021.08.003.
36. **Ostrov** et al., *Highly Specific Sigma Receptor Ligands Exhibit Anti-Viral Properties in SARS-CoV-2 Infected Cells*, *Pathogens*, doi:10.3390/pathogens10111514.
37. **Peacock** et al., *The SARS-CoV-2 variant, Omicron, shows rapid replication in human primary nasal epithelial cultures and efficiently uses the endosomal route of entry*, *bioRxiv*, doi:10.1101/2021.12.31.474653.
38. **Peters**, J., *Comparison of Two Methods to Detect Publication Bias in Meta-analysis*, *JAMA*, doi:10.1001/jama.295.6.676.
39. **Qiao** et al., *Proxalutamide reduces SARS-CoV-2 infection and associated inflammatory response*, *PNAS*, doi:10.1073/pnas.222180912.
40. **Rothstein**, H., *Publication Bias in Meta-Analysis: Prevention, Assessment and Adjustments*, [www.wiley.com/en-ae/Publication+Bias+in+Meta+Analysis:+Prevention,+Assessment+and+Adjustments-p-9780470870143](http://www.wiley.com/en-ae/Publication+Bias+in+Meta+Analysis:+Prevention,+Assessment+and+Adjustments-p-9780470870143).
41. **Rücker** et al., *Arcsine test for publication bias in meta-analyses with binary outcomes*, *Statistics in Medicine*, doi:10.1002/sim.2971.
42. **Said** et al., *The effect of Nigella sativa and vitamin D3 supplementation on the clinical outcome in COVID-19 patients: A randomized controlled clinical trial*, *Frontiers in Pharmacology*, doi:10.3389/fphar.2022.1011522.
43. **Scardua-Silva** et al., *Microstructural brain abnormalities, fatigue, and cognitive dysfunction after mild COVID-19*, *Scientific Reports*, doi:10.1038/s41598-024-52005-7.
44. **Stanley** et al., *Meta-regression approximations to reduce publication selection bias*, *Research Synthesis Methods*, doi:10.1002/jrsm.1095.
45. **Sweeting** et al., *What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data*, *Statistics in Medicine*, doi:10.1002/sim.1761.
46. **Thairu** et al., *A Comparison of Ivermectin and Non Ivermectin Based Regimen for COVID-19 in Abuja: Effects on Virus Clearance, Days-to-discharge and Mortality*, *Journal of Pharmaceutical Research International*, doi:10.9734/jpri/2022/v34i44A36328.
47. **Treanor** et al., *Efficacy and Safety of the Oral Neuraminidase Inhibitor Oseltamivir in Treating Acute Influenza: A Randomized Controlled Trial*, *JAMA*, 2000, 283:8, 1016-1024, doi:10.1001/jama.283.8.1016.
48. **twitter.com**, [twitter.com/FlavioCadegiani/status/1534716400073326592](https://twitter.com/FlavioCadegiani/status/1534716400073326592).

49. **Wan** et al., *Synergistic inhibition effects of andrographolide and baicalin on coronavirus mechanisms by downregulation of ACE2 protein level*, Scientific Reports, doi:10.1038/s41598-024-54722-5.
50. **Willett** et al., *The hyper-transmissible SARS-CoV-2 Omicron variant exhibits significant antigenic change, vaccine escape and a switch in cell entry mechanism*, medRxiv, doi:10.1101/2022.01.03.21268111.
51. **Williams**, T., *Not All Ivermectin Is Created Equal: Comparing The Quality of 11 Different Ivermectin Sources*, Do Your Own Research, [doyourownresearch.substack.com/p/not-all-ivermectin-is-created-equal](https://doyourownresearch.substack.com/p/not-all-ivermectin-is-created-equal).
52. **Xu** et al., *A study of impurities in the repurposed COVID-19 drug hydroxychloroquine sulfate by UHPLC-Q/TOF-MS and LC-SPE-NMR*, Rapid Communications in Mass Spectrometry, doi:10.1002/rcm.9358.
53. **Yang** et al., *SARS-CoV-2 infection causes dopaminergic neuron senescence*, Cell Stem Cell, doi:10.1016/j.stem.2023.12.012.
54. **Zavascki** et al., *Advanced ventilatory support and mortality in hospitalized patients with COVID-19 caused by Gamma (P.1) variant of concern compared to other lineages: cohort study at a reference center in Brazil*, Research Square, doi:10.21203/rs.3.rs-910467/v1.
55. **Zeraatkar** et al., *Consistency of covid-19 trial preprints with published reports and impact for decision making: retrospective review*, BMJ Medicine, doi:10.1136/bmjmed-2022-0003091.
56. **Zhang** et al., *What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes*, JAMA, 80:19, 1690, doi:10.1001/jama.280.19.1690.
57. **Zheng** et al., *Small-molecule antivirals treatment for COVID-19: A systematic review and network meta-analysis*, International Journal of Antimicrobial Agents, doi:10.1016/j.ijantimicag.2024.107096.